

# A Review of Complex Arithmetic

A **complex** value  $C$  has both a **real** and **imaginary** component:


$$a = \text{Re}\{C\} \quad \text{and} \quad b = \text{Im}\{C\}$$

so that we can express this complex value as:

$$C = a + jb$$

where  $j^2 = -1$ .

Just as a **real** value can be expressed as a point on the **real line**, a **complex** value can be expressed as a **point** on the **complex plane**:




The values  $(a,b)$  are a **Cartesian** representation of a point on the complex plane. Recall that we can **alternatively** denote a point on a 2-dimensional plane using **polar** coordinates:

$|C| \doteq$  distance from the origin to the point

$\angle C \doteq \phi_c =$  rotation angle from the horizontal ( $\text{Re}\{C\}$ ) axis

i.e.,



Using our knowledge of **trigonometry**, we can determine the relationship between the Cartesian  $(a,b)$  and polar  $(|C|, \phi_c)$  representations.

From the **Pythagorean** theorem, we find that:

$$|C| = \sqrt{a^2 + b^2}$$

Likewise, from the definition of **sine** (opposite over hypotenuse), we find:

$$\sin \phi_c = \frac{b}{|C|} = \frac{b}{\sqrt{a^2 + b^2}}$$

or, using the definition of **cosine** (adjacent over hypotenuse):

$$\cos \phi_c = \frac{a}{|C|} = \frac{a}{\sqrt{a^2 + b^2}}$$

Combining these results, we can determine the **tangent** (opposite over adjacent) of  $\phi_c$ :

$$\tan \phi_c = \frac{\sin \phi_c}{\cos \phi_c} = \frac{b}{a}$$

Thus, we can write the polar coordinates in terms of the Cartesian coordinates:

$$|C| = \sqrt{a^2 + b^2}$$

$$\phi_c = \tan^{-1}\left(\frac{b}{a}\right) = \cos^{-1}\left(\frac{a}{\sqrt{a^2 + b^2}}\right) = \sin^{-1}\left(\frac{b}{\sqrt{a^2 + b^2}}\right)$$

Likewise, we can use trigonometry to write the **Cartesian** coordinates in terms of the **polar** coordinates.

For example, we can use the definition of *sine* to determine  $b$ :

$$b = |C| \sin \phi_c$$

and the definition of *cosine* to determine  $a$ :

$$a = |C| \cos \phi_c$$

Summarizing:

$$a = |C| \cos \phi_c$$

$$b = |C| \sin \phi_c$$

Note that we can explicitly write the complex value  $C$  in terms of its **magnitude**  $|C|$  and **phase angle**  $\phi_c$ :

$$\begin{aligned} C &= a + jb \\ &= |C| \cos \phi_c + j |C| \sin \phi_c \\ &= |C| (\cos \phi_c + j \sin \phi_c) \end{aligned}$$

*Hey! we can use **Euler's equation** to simplify this further!*

Recall that Euler's equation states:

$$e^{j\phi} = \cos\phi + j \sin\phi$$

so complex value  $C$  is:

$$\begin{aligned} C &= a + jb \\ &= |C|(\cos\phi_c + j \sin\phi_c) \\ &= |C|e^{j\phi_c} \end{aligned}$$

Now we have **two** ways of expressing a complex value  $C$ !

$$C = a + jb \quad \text{and/or} \quad C = |C|e^{j\phi_c}$$

Note that both representations are **equally valid** mathematically—**either one** can be successfully used in complex analysis and computation.

Typically, we find that the **Cartesian** representation is **easiest** to use **if** we are doing **arithmetic** calculations (e.g., addition and subtraction).

For example, if:

$$C_1 = a_1 + j b_1 \quad \text{and} \quad C_2 = a_2 + j b_2$$

then:

$$C_1 + C_2 = (a_1 + a_2) + j(b_1 + b_2)$$

$$C_1 - C_2 = (a_1 - a_2) + j(b_1 - b_2)$$

Conversely, for **geometric** calculations (multiplication and division), it is **easier** to use the **polar** representation:

For example, if:

$$C_1 = |C_1| e^{j\phi_1} \quad \text{and} \quad C_2 = |C_2| e^{j\phi_2}$$

then:

$$\begin{aligned} C_1 C_2 &= |C_1| e^{j\phi_1} |C_2| e^{j\phi_2} \\ &= |C_1| |C_2| e^{j\phi_1} e^{j\phi_2} \\ &= |C_1| |C_2| e^{j(\phi_1 + \phi_2)} \end{aligned}$$

and:

$$\begin{aligned} \frac{C_1}{C_2} &= \frac{|C_1| e^{j\phi_1}}{|C_2| e^{j\phi_2}} \\ &= \frac{|C_1| e^{j\phi_1} e^{-j\phi_2}}{|C_2|} \\ &= \frac{|C_1|}{|C_2|} e^{j(\phi_1 - \phi_2)} \end{aligned}$$

Note in the above calculations we have used the **general** facts:

$$x^y x^z = x^{(y+z)} \quad \text{and} \quad \frac{x^y}{x^z} = x^{(y-z)}$$

Additionally, we note that **powers** and **roots** are most easily accomplished using the **polar** form of  $C$ :

$$\begin{aligned} C^n &= (|C| e^{j\phi_c})^n \\ &= |C|^n (e^{j\phi_c})^n \\ &= |C|^n e^{jn\phi_c} \end{aligned}$$

and

$$\begin{aligned} \sqrt[n]{C} &= (|C| e^{j\phi_c})^{1/n} \\ &= |C|^{1/n} (e^{j\phi_c})^{1/n} \\ &= |C|^{1/n} e^{j(\phi_c/n)} \end{aligned}$$

Therefore:

$$C^2 = (|C| e^{j\phi_c})^2 = |C|^2 e^{j(2\phi_c)}$$

and:

$$\sqrt{C} = (|C| e^{j\phi_c})^{1/2} = \sqrt{|C|} e^{j(\phi_c/2)}$$

Finally, we define the **complex conjugate** ( $C^*$ ) of a complex value  $C$ :

$$\begin{aligned} C^* &\doteq \text{Complex Conjugate of } C \\ &= a - jb \\ &= |C| e^{-j\phi_c} \end{aligned}$$

A **very important** application of the complex conjugate is for determining the **magnitude** of a complex value:

$$|C|^2 = C C^*$$

Typically, the **proof** of this relationship is given as:

$$\begin{aligned} C C^* &= (a + jb)(a - jb) \\ &= a(a - jb) + jb(a - jb) \\ &= a^2 + jab - jba - j^2 b^2 \\ &= a^2 + b^2 \\ &= |C|^2 \end{aligned}$$

However, it is more **easily** shown as:

$$\begin{aligned} C C^* &= (|C| e^{j\phi_c}) (|C| e^{-j\phi_c}) \\ &= |C|^2 e^{j(\phi_c - \phi_c)} \\ &= |C|^2 e^{j0} \\ &= |C|^2 \end{aligned}$$

Another important relationship involving complex conjugate is:

$$\begin{aligned} C + C^* &= (a + jb) + (a - jb) \\ &= (a + a) + j(b - b) \\ &= 2a \end{aligned}$$

Thus, the **sum** of a complex value and its complex conjugate is a purely **real** value.

Additionally, the **difference** of complex value and its complex conjugate results in a purely **imaginary** value:

$$\begin{aligned} C - C^* &= (a + jb) - (a - jb) \\ &= (a - a) + j(b + b) \\ &= j2b \end{aligned}$$

Note from these results we can derive the relationships:

$$a = \text{Re}\{C\} = \frac{C + C^*}{2}$$

$$b = \text{Im}\{C\} = \frac{C - C^*}{j2}$$