

# 4-4 E-field Calculations using Coulomb's Law

Reading Assignment: *pp. 93-98*

Specifically:

1. HO: The Uniform, Infinite Line Charge
2. HO: The Uniform Disk of Charge
3. HO: An Infinite Charge Plane

# The Uniform, Infinite Line Charge


Consider an **infinite** line of charge lying along the  $z$ -axis. The charge density along this line is a **constant** value of  $\rho_\ell$  C/m.

**Q:** *What electric field  $\mathbf{E}(\bar{r})$  is produced by **this** charge distribution?*

**A:** Apply **Coulomb's Law!**

We know that for a **line charge** distribution that:

$$\mathbf{E}(\bar{r}) = \int \frac{\rho_\ell(\bar{r}')}{c} \frac{\bar{r} - \bar{r}'}{|\bar{r} - \bar{r}'|^3} d\ell'$$



**Q:** *Yikes! How do we evaluate **this** integral?*

**A:** Don't panic! **You** know how to evaluate this integral. Let's break up the process into **smaller steps**.

**Step 1:** Determine  $d\ell'$

The differential element  $d\ell'$  is just the **magnitude** of the differential line element we studied in chapter 2 (i.e.,  $d\ell' = |d\vec{\ell}'|$ ). As a result, we can easily integrate over **any** of the seven contours we discussed in chapter 2.

The contour in this problem is one of those! It is a line parallel to the  $z$ -axis, defined as  $x'=0$  and  $y'=0$ . As a result, we use for  $d\ell'$ :

$$d\ell' = |\hat{a}_z dz'| = dz'$$

**Step 2:** Determine the **limits of integration**

This is easy! The line charge is **infinite**. Therefore, we integrate from  $z' = -\infty$  to  $z' = \infty$ .

**Step 3:** Determine the **vector**  $\vec{r} - \vec{r}'$ .

Since for all charge  $x'=0$  and  $y'=0$ , we find:

$$\begin{aligned} \vec{r} - \vec{r}' &= (x\hat{a}_x + y\hat{a}_y + z\hat{a}_z) - (x'\hat{a}_x + y'\hat{a}_y + z'\hat{a}_z) \\ &= (x\hat{a}_x + y\hat{a}_y + z\hat{a}_z) - z'\hat{a}_z \\ &= x\hat{a}_x + y\hat{a}_y + (z - z')\hat{a}_z \end{aligned}$$

**Step 4:** Determine the scalar  $|\bar{r} - \bar{r}'|^3$

Since  $|\bar{r} - \bar{r}'| = \sqrt{x^2 + y^2 + (z - z')^2}$ , we find:

$$|\bar{r} - \bar{r}'|^3 = \left[ x^2 + y^2 + (z - z')^2 \right]^{3/2}$$

**Step 5:** Time to integrate !

$$\begin{aligned} \mathbf{E}(\bar{r}) &= \int_c \frac{\rho_\ell(\bar{r}')}{4\pi\epsilon_0} \frac{\bar{r} - \bar{r}'}{|\bar{r} - \bar{r}'|^3} d\ell' \\ &= \frac{1}{4\pi\epsilon_0} \int_{-\infty}^{\infty} \rho_\ell \frac{x \hat{a}_x + y \hat{a}_y + (z - z') \hat{a}_z}{\left[ x^2 + y^2 + (z - z')^2 \right]^{3/2}} dz' \\ &= \frac{\rho_\ell}{4\pi\epsilon_0} \int_{-\infty}^{\infty} \frac{x \hat{a}_x + y \hat{a}_y + (z - z') \hat{a}_z}{\left[ x^2 + y^2 + (z - z')^2 \right]^{3/2}} dz' \\ &= \frac{\rho_\ell (x \hat{a}_x + y \hat{a}_y)}{4\pi\epsilon_0} \int_{-\infty}^{\infty} \frac{dz'}{\left[ x^2 + y^2 + (z - z')^2 \right]^{3/2}} \\ &\quad + \frac{\rho_\ell \hat{a}_z}{4\pi\epsilon_0} \int_{-\infty}^{\infty} \frac{(z - z') dz'}{\left[ x^2 + y^2 + (z - z')^2 \right]^{3/2}} \\ &= \frac{\rho_\ell (x \hat{a}_x + y \hat{a}_y)}{4\pi\epsilon_0} \frac{2}{x^2 + y^2} + 0 \\ &= \frac{\rho_\ell (x \hat{a}_x + y \hat{a}_y)}{2\pi\epsilon_0 x^2 + y^2} \end{aligned}$$

This result, however, is best expressed in **cylindrical coordinates**:

$$\begin{aligned}\frac{x\hat{a}_x + y\hat{a}_y}{x^2 + y^2} &= \frac{\rho\cos\phi\hat{a}_x + \rho\sin\phi\hat{a}_y}{\rho^2} \\ &= \frac{\cos\phi\hat{a}_x + \sin\phi\hat{a}_y}{\rho}\end{aligned}$$

And with cylindrical **base vectors**:

$$\begin{aligned}\frac{\cos\phi\hat{a}_x + \sin\phi\hat{a}_y}{\rho} &= \frac{1}{\rho}(\cos\phi\hat{a}_x \cdot \hat{a}_\rho + \sin\phi\hat{a}_y \cdot \hat{a}_\rho)\hat{a}_\rho \\ &\quad + \frac{1}{\rho}(\cos\phi\hat{a}_x \cdot \hat{a}_\phi + \sin\phi\hat{a}_y \cdot \hat{a}_\phi)\hat{a}_\phi \\ &\quad + \frac{1}{\rho}(\cos\phi\hat{a}_x \cdot \hat{a}_z + \sin\phi\hat{a}_y \cdot \hat{a}_z)\hat{a}_z \\ &= \frac{1}{\rho}(\cos^2\phi + \sin^2\phi)\hat{a}_\rho \\ &\quad + \frac{1}{\rho}(-\cos\phi\sin\phi + \sin\phi\cos\phi)\hat{a}_\phi \\ &\quad + \frac{1}{\rho}(\cos\phi(0) + \sin\phi(0))\hat{a}_z \\ &= \frac{\hat{a}_\rho}{\rho}\end{aligned}$$

As a result, we can write the **electric field** produced by an **infinite line charge** with constant density  $\rho_\ell$  as:

$$\mathbf{E}(\vec{r}) = \frac{\rho_\ell}{2\pi\epsilon_0} \frac{\hat{a}_\rho}{\rho}$$

Note what this means. Recall unit vector  $\hat{a}_\rho$  is the direction that **points away from** the z-axis. In other words, the electric field produced by the uniform line charge points away from the line charge, just like the electric field produced by a point charge likewise points away from the charge.


It is apparent that the electric field in the static case appears to **diverge** from the location of the charge. And, this is exactly what Maxwell's equations (**Gauss's Law**) says will happen ! i.e.,:

$$\nabla \cdot \mathbf{E}(\vec{r}) = \frac{\rho_v(\vec{r})}{\epsilon_0}$$

Note the **magnitude** of the electric field is proportional to  $1/\rho$ , therefore the electric field **diminishes** as we get further from the line charge. Note however, the electric field does not diminish as **quickly** as that generated by a point charge. Recall in that case, the magnitude of the electric field diminishes as  $1/r^2$ .

# The Uniform Disk of Charge

Consider a **disk** radius  $a$ , centered at the origin, and lying entirely on the  $z=0$  plane.



This disk contains **surface charge**, with density of  $\rho_s$  C/m<sup>2</sup>. This density is **uniform** across the disk.

Let's find the **electric field** generated by this charge disk!

From **Coulomb's Law**, we know:

$$\mathbf{E}(\bar{r}) = \iint_S \frac{\rho_s(\bar{r}')}{4\pi\epsilon_0} \frac{\bar{r} - \bar{r}'}{|\bar{r} - \bar{r}'|^3} ds'$$

**Step 1:** Determine  $ds'$

This disk can be described by the equation  $z' = 0$ . That is, every point on the disk has a coordinate value  $z'$  that is equal to zero.

This is **one** of the surfaces we examined in chapter 2. The **differential surface element** for that surface, you recall, is:

$$ds' = ds_z = \rho' d\rho' d\phi'$$

**Step 2:** Determine the **limits of integration**.

Note over the surface of the disk,  $\rho'$  changes from 0 to radius  $a$ , and  $\phi'$  changes from 0 to  $2\pi$ . Therefore:

$$0 < \rho' < a \quad 0 < \phi' < 2\pi$$

**Step 3:** Determine vector  $\bar{r} - \bar{r}'$ .

We know that  $z' = 0$  for all charge, therefore we can write:

$$\begin{aligned} \bar{r} - \bar{r}' &= (x\hat{a}_x + y\hat{a}_y + z\hat{a}_z) - (x'\hat{a}_x + y'\hat{a}_y + z'\hat{a}_z) \\ &= (x\hat{a}_x + y\hat{a}_y + z\hat{a}_z) - (x'\hat{a}_x + y'\hat{a}_y) \\ &= (x - x')\hat{a}_x + (y - y')\hat{a}_y + z\hat{a}_z \end{aligned}$$

Since the primed coordinates in  $ds'$  are expressed in **cylindrical** coordinates, we convert the coordinates to get:



$$\begin{aligned}
 \bar{r} - \bar{r}' &= (x \hat{a}_x + y \hat{a}_y + z \hat{a}_z) - (x' \hat{a}_x + y' \hat{a}_y) \\
 &= (x - x') \hat{a}_x + (y - y') \hat{a}_y + z \hat{a}_z \\
 &= (x - \rho' \cos \phi') \hat{a}_x + (y - \rho' \sin \phi') \hat{a}_y + z \hat{a}_z
 \end{aligned}$$

**Step 4:** Determine  $|\bar{r} - \bar{r}'|^3$

We find that:

$$|\bar{r} - \bar{r}'|^3 = \left[ (x - \rho' \cos \phi')^2 + (y - \rho' \sin \phi')^2 + z^2 \right]^{3/2}$$

**Step 5:** Time to integrate !

$$\begin{aligned}
 \mathbf{E}(\bar{r}) &= \iint_S \frac{\rho_s(\bar{r}')}{4\pi\epsilon_0} \frac{\bar{r} - \bar{r}'}{|\bar{r} - \bar{r}'|^3} ds' \\
 &= \frac{\rho_s}{4\pi\epsilon_0} \int_0^{2\pi} \int_0^a \frac{(x - \rho' \cos \phi') \hat{a}_x + (y - \rho' \sin \phi') \hat{a}_y + z \hat{a}_z}{\left[ (x - \rho' \cos \phi')^2 + (y - \rho' \sin \phi')^2 + z^2 \right]^{3/2}} \rho' d\rho' d\phi'
 \end{aligned}$$

Yikes! What a **mess!** To **simplify** our integration let's determine the electric field  $\mathbf{E}(\bar{r})$  along the **z-axis** only. In other words, set  $x = 0$  and  $y = 0$ .

$$\begin{aligned}
\mathbf{E}(x=0, y=0, z) &= \iint_S \frac{\rho_s(\vec{r}')}{4\pi\epsilon_0} \frac{\vec{r} - \vec{r}'}{|\vec{r} - \vec{r}'|^3} ds' \\
&= \frac{\rho_s}{4\pi\epsilon_0} \int_0^{2\pi} \int_0^a \frac{(0 - \rho' \cos\phi') \hat{a}_x + (0 - \rho' \sin\phi') \hat{a}_y - z \hat{a}_z}{\left[ (0 - \rho' \cos\phi')^2 + (0 - \rho' \sin\phi')^2 + z^2 \right]^{3/2}} \rho' d\rho' d\phi' \\
&= \frac{-\rho_s}{4\pi\epsilon_0} \int_0^{2\pi} \int_0^a \frac{(\rho' \cos\phi') \hat{a}_x + (\rho' \sin\phi') \hat{a}_y - z \hat{a}_z}{\left[ \rho'^2 + z^2 \right]^{3/2}} \rho' d\rho' d\phi' \\
&= \frac{\rho_s}{4\pi\epsilon_0} \hat{a}_x \int_0^{2\pi} \int_0^a \frac{(\rho' \cos\phi') \rho' d\rho' d\phi'}{\left[ \rho'^2 + z^2 \right]^{3/2}} \\
&\quad + \frac{-\rho_s}{4\pi\epsilon_0} \hat{a}_y \int_0^{2\pi} \int_0^a \frac{(\rho' \sin\phi') \rho' d\rho' d\phi'}{\left[ \rho'^2 + z^2 \right]^{3/2}} \\
&\quad + \frac{-\rho_s}{4\pi\epsilon_0} \hat{a}_z \int_0^{2\pi} \int_0^a \frac{z \rho' d\rho' d\phi'}{\left[ \rho'^2 + z^2 \right]^{3/2}}
\end{aligned}$$


Note that since:

$$\int_0^{2\pi} \sin\phi d\phi = 0 = \int_0^{2\pi} \cos\phi d\phi$$

The first two terms ( $E_x$  and  $E_y$ ) are equal to zero. Integrating the last term, we get:

$$\mathbf{E}(x=0, y=0, z) = \begin{cases} \frac{\rho_s}{2\epsilon_0} \hat{a}_z \left[ 1 - \frac{z}{\sqrt{z^2 + a^2}} \right] & \text{if } z > 0 \\ \frac{\rho_s}{2\epsilon_0} \hat{a}_z \left[ -1 - \frac{z}{\sqrt{z^2 + a^2}} \right] & \text{if } z < 0 \end{cases}$$

From this expression, we can conclude **two** things. The first is that **above** the disk ( $z > 0$ ), the electric field points in the direction  $\hat{a}_z$ , and below the disk ( $z < 0$ ), it points in the direction  $-\hat{a}_z$ .



What a surprise (not)! The electric field **points away** from the charge. It appears to be **diverging** from the charged disk (as predicted by Gauss's Law).

Likewise, it is evident that as we move further and **further from** the disk, the electric field will **diminish**. In fact, as distance  $z$  goes to **infinity**, the magnitude of the electric field approaches **zero**. This of course is similar to the **point** or **line** charge; as we move an infinite distance away, the electric field diminishes to **nothing**.

# An Infinite Charge Plane

Say that we have a **very large** charge disk. So large, in fact, that its radius  $a$  approaches **infinity** !

**Q:** *What electric field is created by this infinite plane?*

**A:** We **already** know! Just evaluate the charge disk solution for the case where the disk **radius**  $a$  is **infinity**.

In other words:

$$\lim_{a \rightarrow \infty} \mathbf{E}(x=0, y=0, z) = \begin{cases} \hat{a}_z \frac{\rho_s}{2\epsilon_0} \left[ 1 - \frac{z}{\sqrt{z^2 + a^2}} \right] & \text{if } z > 0 \\ \hat{a}_z \frac{\rho_s}{2\epsilon_0} \left[ -1 - \frac{z}{\sqrt{z^2 + a^2}} \right] & \text{if } z < 0 \end{cases}$$


$$= \begin{cases} \frac{\rho_s}{2\epsilon_0} \hat{a}_z & \text{if } z > 0 \\ -\frac{\rho_s}{2\epsilon_0} \hat{a}_z & \text{if } z < 0 \end{cases}$$

Therefore, the electric field produced by an **infinite charge plane**, with surface charge density  $\rho_s$ , is:

$$\mathbf{E}(\vec{r}) = \begin{cases} \frac{\rho_s}{2\epsilon_0} \hat{a}_z & \text{if } z > 0 \\ -\frac{\rho_s}{2\epsilon_0} \hat{a}_z & \text{if } z < 0 \end{cases}$$

Think about what **this** says!

- \* First, we note that the electric field **points away** from the plane if  $\rho_s$  is positive, and toward the plane if  $\rho_s$  is negative.
- \* Second, we notice that the magnitude of the electric field is a **constant**—the magnitude is **independent** of the distance from the infinite plane!



The reason for this result is, that no matter how far you are (i.e.,  $|z|$ ) from the infinite charge plane, you remain **infinitely close** to plane, when **compared** to its radius  $a$ .

We will find these results are useful when we study the behavior of a parallel plate **capacitor**.